

Resistenzlage Klinikum am Urban 2019

Zeitraum: 01.01.2019 bis 31.12.2019

Anteil der resistent getesteten Erreger in Prozent (%)

	Anzahl Testungen	Penicillin	Oxacillin	Ampicillin	Ampicillin/Sulbactam	Piperacillin	Piperacillin/Tazobactam	Cefazolin	Cefuroxim	Cefotaxim	Ceftriaxon	Ceftazidim	Aztreonam	Imipenem	Meropenem	Gentamicin	Gentamicin HL	Tobramycin	Ciprofloxacin	Moxifloxacin	Tigecyclin	Co-Trimoxazol	Colistin	Vancomycin	Teicoplanin	Erythromycin	Clindamycin	Linezolid	Rifampicin	Fosfomycin	Daptomycin	Fusidinsäure
Staph. aureus* ¹	551	n.d.	11	n.d.	11	n.d.	11	11	11	11	11	n.d.	n.d.	11	11	1	n.d.	1	20	17	0	3	n.d.	0	n.d.	20	18	0	1	2	0	4
Koagulase neg. Staphylokokken* ¹	793	n.d.	53	n.d.	53	n.d.	53	53	53	53	53	n.d.	n.d.	53	53	22	n.d.	23	36	33	1	21	n.d.	0	n.d.	56	39	0	3	41	1	33
Enterokokken* ²	946	n.d.	n.d.	21	21	21	21	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	21	n.d.	n.d.	n.d.	n.d.	100	n.d.	0	100	n.d.	7	4	n.d.	n.d.	0	n.d.	n.d.	n.d.	n.d.
E. coli	1341	n.d.	n.d.	50	47	50	18	n.d.	19	15	15	n.d.	n.d.	0	0	5	n.d.	n.d.	24	31	0	24	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Klebsiella	519	n.d.	n.d.	100	24	99	18	n.d.	14	10	10	n.d.	n.d.	0	0	3	n.d.	n.d.	13	17	3	12	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Enterobacter	201	n.d.	n.d.	100	99	25	22	n.d.	100	20	21	21	n.d.	1	1	1	n.d.	n.d.	3	8	0	4	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Proteus	349	n.d.	n.d.	50	34	26	3	n.d.	14	3	3	3	n.d.	8	0	8	n.d.	n.d.	13	54	n.d.	23	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
P. aeruginosa	242	n.d.	n.d.	n.d.	n.d.	19	18	n.d.	n.d.	n.d.	n.d.	12	16	19	11	4	n.d.	2	15	n.d.	n.d.	n.d.	0	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.

Resistenz < 10%

Resistenz 10-49%

Resistenz >=50%

n.d. nicht getestet

*1: Ampicillin/Sulbactam, Piperacillin/Tazobactam, Cefazolin, Cefuroxim, Cefotaxim, Ceftriaxon, Imipenem, Meropenem sind von Oxacillin/Cefoxitin abgeleitet

*2: Ampicillin/Sulbactam, Piperacillin, Piperacillin/Tazobactam sind von Ampicillin abgeleitet