

# Resistenzlage Klinikum Kaulsdorf 2019

Zeitraum: 01.01.2019 bis 31.12.2019

Anteil der resistent getesteten Erreger in Prozent (%)

	Anzahl Testungen	Penicillin	Oxacillin	Ampicillin	Ampicillin/Sulbactam	Piperacillin	Piperacillin/Tazobactam	Cefazolin	Cefuroxim	Cefotaxim	Ceftriaxon	Ceftazidim	Aztreonam	Imipenem	Meropenem	Gentamicin	Gentamicin HL	Tobramycin	Ciprofloxacin	Moxifloxacin	Tigecyclin	Co-Trimoxazol	Colistin	Vancomycin	Teicoplanin	Erythromycin	Clindamycin	Linezolid	Rifampicin	Fosfomycin	Daptomycin	Fusidinsäure
Staph. aureus* <sup>1</sup>	211	n.d.	7	n.d.	7	n.d.	7	7	7	7	7	n.d.	n.d.	7	7	0	n.d.	1	18	15	0	1	n.d.	0	n.d.	13	10	0	0	0	0	0
Koagulase neg. Staphylokokken* <sup>1</sup>	269	n.d.	51	n.d.	51	n.d.	51	51	51	51	51	n.d.	n.d.	51	51	22	n.d.	22	38	31	0	13	n.d.	0	n.d.	55	45	0	5	47	1	33
Enterokokken* <sup>2</sup>	331	n.d.	n.d.	23	23	23	23	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	23	n.d.	n.d.	n.d.	n.d.	100	n.d.	0	99	n.d.	8	4	n.d.	n.d.	0	n.d.	n.d.	n.d.	n.d.
E. coli	618	n.d.	n.d.	49	43	48	16	n.d.	19	13	13	n.d.	n.d.	0	0	4	n.d.	n.d.	19	26	0	22	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Klebsiella	186	n.d.	n.d.	100	22	100	16	n.d.	15	8	8	n.d.	n.d.	1	1	2	n.d.	n.d.	9	15	7	9	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Enterobacter	56	n.d.	n.d.	100	100	29	29	n.d.	100	29	29	29	n.d.	0	0	0	n.d.	n.d.	2	9	0	2	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Proteus	147	n.d.	n.d.	43	25	23	6	n.d.	11	5	5	5	n.d.	7	0	7	n.d.	n.d.	12	54	n.d.	25	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
P. aeruginosa	91	n.d.	n.d.	n.d.	n.d.	9	8	n.d.	n.d.	n.d.	n.d.	7	11	8	3	1	n.d.	0	9	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.

Resistenz < 10%

Resistenz 10-49%

Resistenz >=50%

n.d. nicht getestet

\*1: Ampicillin/Sulbactam, Piperacillin/Tazobactam, Cefazolin, Cefuroxim, Cefotaxim, Ceftriaxon, Imipenem, Meropenem sind von Oxacillin/Cefoxitin abgeleitet

\*2: Ampicillin/Sulbactam, Piperacillin, Piperacillin/Tazobactam sind von Ampicillin abgeleitet