

Resistenzlage Vivantes-Kliniken 2019

Zeitraum: 01.01.2019 bis 31.12.2019

Anteil der resistent getesteten Erreger in Prozent (%)

	Anzahl Testungen	Oxacillin	Ampicillin	Ampicillin/Sulbactam	Piperacillin	Piperacillin/Tazobactam	Cefazolin	Cefuroxim	Cefotaxim	Ceftriaxon	Ceftazidim	Aztreonam	Imipenem	Meropenem	Gentamicin	Gentamicin HL	Tobramycin	Ciprofloxacin	Moxifloxacin	Tigecyclin	Co-Trimoxazol	Colistin	Vancomycin	Teicoplanin	Erythromycin	Clindamycin	Linezolid	Rifampicin	Fosfomycin	Daptomycin	Fusidinsäure
Staph. aureus* ¹	5833	10	n.d.	10	n.d.	10	10	10	10	10	n.d.	n.d.	10	10	1	n.d.	1	19	17	0	2	n.d.	0	n.d.	18	16	0	0	1	0	3
Koagulase neg. Staphylokokken* ¹	7332	53	n.d.	53	n.d.	53	53	53	53	53	n.d.	n.d.	53	53	25	n.d.	26	40	37	0	21	n.d.	0	n.d.	57	40	0	4	42	0	32
Enterokokken* ²	7890	n.d.	21	21	21	21	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	21	n.d.	n.d.	n.d.	n.d.	100	n.d.	0	99	n.d.	7	4	n.d.	n.d.	0	n.d.	n.d.	n.d.	n.d.
E. coli	11503	n.d.	48	44	47	16	n.d.	17	12	12	n.d.	n.d.	0	0	6	n.d.	n.d.	20	26	0	23	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Klebsiella	4485	n.d.	100	24	100	18	n.d.	15	9	9	n.d.	n.d.	0	0	3	n.d.	n.d.	10	15	4	9	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Enterobacter	1947	n.d.	99	99	27	24	n.d.	100	23	24	23	n.d.	1	0	1	n.d.	n.d.	3	8	3	3	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Proteus	3475	n.d.	50	32	27	3	n.d.	16	4	4	4	n.d.	8	0	7	n.d.	n.d.	13	53	n.d.	24	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
P. aeruginosa	2549	n.d.	n.d.	n.d.	16	14	n.d.	n.d.	n.d.	n.d.	10	13	16	7	3	n.d.	2	12	n.d.	n.d.	n.d.	6	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.

Resistenz < 10%
Resistenz 10-49%
Resistenz >=50%
n.d. nicht getestet

*1: Ampicillin/Sulbactam, Piperacillin/Tazobactam, Cefazolin, Cefuroxim, Cefotaxim, Ceftriaxon, Imipenem, Meropenem sind von Oxacillin/Cefoxitin abgeleitet

*2: Ampicillin/Sulbactam, Piperacillin, Piperacillin/Tazobactam sind von Ampicillin abgeleitet