

Resistenzlage Wenckebach-Klinikum 2019

Zeitraum: 01.01.2019 bis 31.12.2019

Anteil der resistent getesteten Erreger in Prozent (%)

	Anzahl Testungen	Penicillin	Oxacillin	Ampicillin	Ampicillin/Sulbactam	Piperacillin	Piperacillin/Tazobactam	Cefazolin	Cefuroxim	Cefotaxim	Ceftriaxon	Ceftazidim	Aztreonam	Imipenem	Meropenem	Gentamicin	Gentamicin HL	Tobramycin	Ciprofloxacin	Moxifloxacin	Tigecyclin	Co-Trimoxazol	Colistin	Vancomycin	Teicoplanin	Erythromycin	Clindamycin	Linezolid	Rifampicin	Fosfomycin	Daptomycin	Fusidinsäure
Staph. aureus ^{*1}	212	n.d.	11	n.d.	11	n.d.	11	11	11	11	11	n.d.	n.d.	11	11	3	n.d.	3	26	24	0	1	n.d.	0	n.d.	18	16	0	0	3	0	2
Koagulase neg. Staphylokokken ^{*1}	327	n.d.	53	n.d.	53	n.d.	53	53	53	53	53	n.d.	n.d.	53	53	24	n.d.	24	36	36	0	20	n.d.	0	0	53	38	0	2	43	0	32
Enterokokken ^{*2}	429	n.d.	n.d.	22	22	22	22	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	22	n.d.	n.d.	n.d.	n.d.	100	n.d.	0	100	n.d.	5	3	n.d.	n.d.	0	n.d.	n.d.	n.d.	n.d.
E. coli	742	n.d.	n.d.	47	41	44	12	n.d.	17	10	10	n.d.	n.d.	0	0	5	n.d.	n.d.	19	25	0	19	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Klebsiella	241	n.d.	n.d.	100	20	100	15	n.d.	11	6	6	n.d.	n.d.	0	0	3	n.d.	n.d.	7	12	1	7	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Enterobacter	79	n.d.	n.d.	100	100	30	27	n.d.	100	26	26	26	n.d.	0	0	0	n.d.	n.d.	1	3	0	6	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Proteus	212	n.d.	n.d.	49	31	26	3	n.d.	18	3	3	3	n.d.	8	0	8	n.d.	n.d.	12	53	n.d.	20	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
P. aeruginosa	111	n.d.	n.d.	n.d.	n.d.	13	9	n.d.	n.d.	n.d.	n.d.	13	3	4	0	9	n.d.	0	9	n.d.	n.d.	n.d.	0	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.

Resistenz < 10%

Resistenz 10-49%

Resistenz >=50%

n.d. nicht getestet

*1: Ampicillin/Sulbactam, Piperacillin/Tazobactam, Cefazolin, Cefuroxim, Cefotaxim, Ceftriaxon, Imipenem, Meropenem sind von Oxacillin/Cefoxitin abgeleitet

*2: Ampicillin/Sulbactam, Piperacillin, Piperacillin/Tazobactam sind von Ampicillin abgeleitet